



# Emergency Care for **CHOKING**


## CONSCIOUS VICTIM

If victim **CAN** breathe,  
cough or make sounds,  
**DO NOT INTERFERE.**



**Give quick upward thrusts above the belly button and below the ribs until object is forced out, victim can breathe again, or victim becomes unconscious**

If victim **CANNOT** breathe,  
cough or make sounds,  
ask if you can help.



## UNCONSCIOUS VICTIM

Send someone to call 911 and get the Automated External Defibrillator (AED).  
**IF YOU ARE ALONE**, perform 5 sets of 30 compressions and 2 breaths before leaving to call 911. Follow these steps.



**1**  
Give 30 compressions pushing down **AT LEAST 2 inches** on the center of the chest. Place one hand on top of the other. Push hard.



**2**  
Open the airway and check the mouth for objects. Remove the obstructing object only if you see it.



**3**  
With the airway open, attempt to give **TWO** breaths. If unsuccessful, return to compressions.

**Repeat steps 1, 2 and 3 until victim starts breathing or until emergency medical help arrives.**

Illinois Department of Public Health  
Emergency Medical Systems and Highway Safety  
422 S. 5th St., Third Floor  
Springfield, IL 62701 • 217-785-2080

Standards for CPR and ECC are consistent with American Heart Association recommendations.

- Have someone call for an ambulance, rescue squad or EMS.
- **DO NOT PRACTICE ON PEOPLE.** Abdominal thrusts may cause injury.
- Use back blows and chest thrust on infants. Use chest thrust on pregnant women and obese victims.
- For children 1 to 8 years of age, compress at the depth of approximately 2 inches.
- Learn to perform emergency care for choking and cardiopulmonary resuscitation (CPR).
- For CPR training information, call your local American Heart Association or American Red Cross chapter.